

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified Applicators or persons under their direct supervision and only for those uses covered by the certified Applicators certification.

CPF 4E

Chlorpyrifos Agricultural Insecticide

For control of listed insects infesting certain field, fruit, nut, and vegetable crops and wheat.

Active Ingredient:

Chlorpyrifos:

O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate 45.0%

Inert Ingredients :..... 55.0%

Total100.0%

Contains petroleum distillate

Contains 4 pounds of Chlorpyrifos per gallon.

EPA Reg. No. 83222-20

EPA Establishment No. 53883-TX-002

44616-MO-001

39578-TX-001

**KEEP OUT OF REACH OF CHILDREN
WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID (Organophosphate Insecticide)	
If inhaled:	<ul style="list-style-type: none"> Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER (Organophosphate Insecticide)	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information call: 1-800-222-1222	

NOTE TO PHYSICIAN

Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell Cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration. Note: Contains Petroleum Distillate -vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING. May Be Fatal If Swallowed. Harmful If Absorbed Through The Skin. Causes Moderate Eye And Skin Irritation. Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Materials that are chemical-resistant to this product are Barrier Laminate and Viton \geq 14 mils.

All mixers, loaders, other applicators and other handlers must wear:

- coveralls over long-sleeved shirt and long pants;
- chemical-resistant gloves;
- chemical-resistant apron when mixing or loading or exposed to the concentrate;
- chemical resistant footwear plus socks;
- chemical-resistant headgear for overhead exposures;
- a NIOSH-approved dust/mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any R, P or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When applicators use closed equipment in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, small mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Cover or incorporate spills. Do not contaminate water when disposing of equipment wash water or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Combustible. Do not use or store near heat or open flame. Do not cut or weld container.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

This product cannot be reformulated or repackaged into other end-use products.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer at the end of this label.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the required restricted entry interval (REI). The required REI's are: For all crops: 24 hours, unless specifically otherwise noted below

- Fruit trees (including fig): 4 days
- Citrus trees: 5 days
- Cauliflower: 3 days

Also see specific Use Directions under Approved Crops Section of this label

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls over short-sleeved shirt and short pants;
- chemical-resistant gloves made out of water proof material;
- chemical-resistant footwear plus socks;
- chemical-resistant headgear for overhead exposure.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

Product Information

CPF 4E insecticide forms an emulsion when diluted with water and is suitable for use in all conventional spray equipment. Consult your State Experiment Station or State Extension Service for proper timing of applications.

Mixing Directions

To prepare the spray, add a portion of the required amount of water to the spray tank and with the spray tank agitator operating add the CPF4E. Complete filling the tank with the balance of water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

CPF 4E may also be used in tank mixtures with certain herbicides and/or with non-pressure fertilizer solutions as recommended under specific crop use directions. Prepare tank mixtures in the same manner as recommended above for use of CPF 4E alone. When tank mixtures of CPF 4E and herbicides are involved, add wettable powders first, flowables second, and emulsifiable concentrates last. Where a fertilizer solution is involved, it is strongly recommended that a fertilizer pesticide compatibility agent such as Unite or Complex be used. Maintain constant agitation during both mixing and application to ensure uniformity of the spray mixture. Do not allow spray mixtures to stand overnight.

Note: Test compatibility of the intended tank mixture before adding CPF 4E to the spray or mix tank. Add proportionate amounts of each ingredient to a pint or quart jar, cap, shake, and let set 15 minutes. Formation of precipitates that do not readily re-disperse indicates an incompatible mixture that should not be used.

Sprinkler Irrigation

CPF 4E may be applied by sprinkler irrigation for the following crop uses: alfalfa, citrus, almond and walnut orchard floors, field corn, mint, sweet corn, cotton, cranberries, sorghum, and soybeans. Do not apply this product through any type of irrigation system for crops not listed above.

Sprinkler Irrigation Use Directions

The following use directions are to be followed when CPF 4E is applied through sprinkler irrigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injector with soap and water. Determine the amount of insecticide needed to cover the desired acreage. Pump the required CPF 4E into a steel tank, start mechanical or hydraulic agitation, and add in order the non-emulsifiable oil and/or water. Continually agitate the mixture containing CPF 4E. Set the sprinkler system to deliver the desired inches of water per acre. Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injector system according to Step 14 in the "Special Use Precautions" Section of this label. The mixture containing CPF 4E must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

Sprinkler Irrigation Use Precautions

The following use precautions will result in a safe and successful application of mixtures containing CPF 4E. See the use sections for the individual crops for further application information.

1. Apply this product only through sprinkler irrigation systems including center pivot, lateral move and tow, side (wheel) roll, traveler, big gun, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
3. If you have questions about calibration, you should contact state extension service specialists, equipment manufacturers, or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. The metering pump must provide a greater pressure than that of the irrigation system at the point of injection. The pump must meet Section 675 for Electrically Driven or Controlled Irrigation Machines NEC 70 and must contain Viton or Teflon seals.
12. To insure uniform mixing of the insecticide into the water line, inject the mixture through a nozzle placed in the fertilizer injection port or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. It is suggested that the injection point be higher than the insecticide tank to prevent siphoning.
13. The steel tank holding the insecticide mixture should be large enough to allow the system to complete a revolution with 1 filling. It should be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector pump.
14. In order to calibrate the irrigation system and injector to apply the mixture containing CPF 4E, determine the following: 1) Calculate the number of acres irrigated by the system; 2) Set the irrigation rate and determine the number of minutes for the system to cover the intended treatment area; 3) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes to cover the treatment area. This value equals the gallons per minute to milliliters or ounces per minute. Calibrate the injector pump with the system in operation at the desired irrigation rate. It is suggested that the injector pump be calibrated at least twice before operation, and the system should be monitored during operation.
15. Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application, if they irrigate non-target areas.
16. Do not allow irrigation water to collect or runoff and pose a hazard to livestock, wells, or adjoining crops.
17. Follow all Worker protection standards (WPS) and reentry restriction on this label.
18. Do not apply through sprinkler systems which deliver a low coefficient of uniformity such as certain water drive units.
19. Do not apply within 25 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries or commercial fish ponds.
20. Applications of CPF 4E to labeled crops growing under environmental conditions that combine high elevation, high heat, or bright sunlight may have greater potential for injury. Application techniques that may reduce possible harmful effects include: spraying later in the day when air temperatures are cooling, using higher amounts of water with the application, avoiding tank mix combinations which require the use of additional adjuvant or surfactants, and using the lower end of the rate range for a particular insect.

Application Precautions and Restrictions

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application-

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks, and recreation areas, non target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

For ground boom applications do not apply within 25 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries or commercial fish ponds. Apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer. Use fine or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.

For orchard/vineyard airblast applications do not apply within 50 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries or commercial fish ponds. Direct spray above trees/vines and turn off outward pointing nozzles at row ends and outer rows. Apply only when wind speed is 3-10mph at the application site as measured by a anemometer outside of the orchard/vineyard on the upwind side.

For aerial applications, do not apply within 150 feet of rivers, natural ponds lakes, streams, reservoirs, marshes, estuaries or commercial fish ponds. The boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3-10 mph as measured by an anemometer. Use fine or courser spray according to ASAE572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or the crop canopy.

For overhead chemigation do not apply within 25 feet of rivers, natural ponds. lakes, streams, reservoirs, marshes, estuaries or commercial fish ponds. Apply only when wind speed is 10 mph or less.

The applicator also must use all other measures necessary to control drift.

Approved Crops

Alfalfa

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Foliar Applications:

Use CPF 4E to control the following pests at the dosages indicated by application as a broadcast, foliar spray:

Pests	Dosage of CPF 4E
corn rootworm adults (spotted cucumber beetle) grasshoppers leafhoppers	1/2 - 1 pt/acre
alfalfa blotch leafminer alfalfa caterpillar alfalfa weevil larvae and adults armyworms blue alfalfa aphid cutworms Egyptian alfalfa weevil larvae and adults pea aphid plant bugs spittlebugs spotted alfalfa aphid (suppression) [not registered for use in California)	1 - 2 pt/acre

Additional Geographical Rate Instructions: Use higher rates within the specified rate range to control spotted alfalfa aphid in Nevada. Stubble spray may be applied to control leafhopper in the Northeast.

Mix the required dosage with enough water to ensure thorough coverage of crop foliage and apply using aerial (fixed-wing or helicopter) or power operated ground spray equipment.

Control may be reduced at low spray volumes under high temperature and wind conditions. Treat when field counts or crop injury indicates that damaging pest populations are developing or present; however, do not apply more than

once per crop cutting. Some reduction in insect control may be evident under excessively cool conditions. For Egyptian alfalfa weevil control in California, apply the specified dosage in a minimum of 5 gallons of water per acre when larvae are actively feeding and populations reach 15 to 20 larvae per 180° sweep with a 15-inch diameter net.

Do not tank mix CPF 4E with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination non-injurious under your current conditions of use. Some phytotoxic symptoms may be observed on young, tender, rapidly growing alfalfa when treated with CPF 4E. Alfalfa will outgrow the symptoms and no yield loss should be expected.

Aerial Applications: For aerial application use 2 to 5 gallons of water per acre. For best coverage when using ground application, a minimum of 20 gallons of water per acre with hollow cone nozzles is recommended.

Sprinkler Irrigation: CPF 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the recommended rate of CPF 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" for further information.

Precautions: This product is highly toxic to bees exposed to direct treatment on alfalfa. Do not apply if nearby bees are clustered outside of hives and bees are foraging. Protective information may be obtained from your Agricultural Extension Service.

Restrictions: Do not cut or graze treated alfalfa within 7 days after application of 1/2 pint of CPF 4E per acre, within 14 days after application of 1 pint per acre, or within 21 days after application of rates above 1 pint per acre. Do not make more than 4 applications per year or apply more than once per crop cutting. Maximum single application rate is 1 lb ai chlorpyrifos (2 pints CPF 4E) per acre. Do not make a second application of CPF 4E or any other product containing chlorpyrifos within 10 days of the first application.

Asparagus

[For use only in the Midwest and Pacific Northwest (AZ, CA, ID, IN, IA, KS, MI, MN, MO, NE, ND, OH, OR, SD, WA, WI)]

<p>Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless Personal Protective Equipment (PPE) required for early entry is worn.</p>

Use CPF 4E to control cutworms, asparagus aphids, and asparagus beetles by application at the rate of 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Mix the specified dosage in sufficient water to ensure thorough coverage of treated plants and apply as a ground, broadcast, foliar spray. For cutworms, it is preferable to apply CPF 4E when the soil is moist and worms are active on or near the soil surface. Applications may be made during the fern stage for control of asparagus beetles and asparagus aphids when field counts or crop injury indicates that damaging pest populations are developing or present.

Restrictions: Do not make more than 1 preharvest application per season or apply within 1 day of harvest. Do not make more than 2 postharvest applications during the fern stage. Based on available residue data, the use of CPF 4E on asparagus is limited to the Midwest and Pacific Northwest (AZ, CA, ID, IN, IA, KS, MI, MN, MO, NE, ND, OH, OR, SD, WA, WI). Maximum single application rate is 1 lb ai chlorpyrifos (2 pints CPF 4E) per acre. Do not make a second application of CPF 4E or any other product containing chlorpyrifos within 10 days of the first application.

Christmas Trees (Plantations Only)

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless Personal Protective Equipment (PPE) required for early entry is worn.

Unless otherwise indicated, apply as a foliar spray using power-operated ground equipment. Use CPF 4E at the rate indicated to control the following insects on the tree varieties listed.

Do not allow livestock to graze in treated areas.

Nurseries and Plantation Crops

Tree Variety	Insects Controlled	Dosage of CPF 4E	Remarks
balsam fir blue spruce concolor fir Douglas fir Eastern white pine Fraser fir grand fir noble fir Scotch pine white spruce	Ants (excluding fire, harvester, carpenter, and pharaoh ants) aphids adelgids (cooley, Eastern spruce gal) European pine sawfly European pine Shoot moth grasshoppers gypsy moth mites ¹ (European red spider, two spotted spider) [except in WA or OR] pales weevil (adult) pine needle midge Douglas fir needle midge pine spittlebug plant bugs spittlebugs spruce budworm spruce needle-miner scale ² (pine needle) (pine tortoise) (spruce bud) (black Pine) (striped pine)	1 qt/acre	Do not treat plants under extreme heat and drought stress. Apply to foliage in sufficient water to ensure adequate coverage. ¹ For effective control of adult spider mites if large numbers of eggs are present, apply a second spray 7 to 10 days after initial treatment to control newly hatched nymphs. ² For scale control apply when scale crawlers are active.
	pales weevil (Beetle)	3 gt/100 gal	Apply as a cut stump drench.

Citrus Fruits (Calamondin, chironja, citrus citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin (tangerine), pummelo, satsuma mandarin, sour orange, sweet orange, tangelo, tangor)
(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E at the rates indicated according to the designated geographic area and pest. Use the lower rates in the rate range for light infestations and the higher rates in the rate range for heavier infestations.

Petroleum spray oil approved for use on citrus trees may be added to dilute spray mixtures only at a rate of up to 1.8 gallons per 100 gallons of water to improve control of aphids, mealybugs, scale insects, and thrips. Treat when insects become a problem or in accordance with the local spray schedule recommended by your State Extension Service Specialist.

Approved Applications for Citrus by Geographical Area

Geographic Location	Pest	Dosage of CPF 4E (pt/acre)	Spray Volume (gal/acre)
California Arizona	aphids katydids Lepidopterous larvae Avocado leafroller cutworms Fruit tree leafroller Orange tortrix Western tussock moth	2 - 8	ground: 100-750 aerial: Min. of 15
California Arizona	(Nips (suppression) Mealybugs	6 - 8	100-750
California Arizona	Scale insects black scale brown scale	4 - 8	100-2400
California*	Scale California red scale	8 - 12	100-2400

Restriction: Do not use a spray concentration of CPF 4E of less than 1/2 pt/100 gal of total volume.

<p>* Additional Precautions for California and Arizona: Maximum of 8 pints per acre except in Fresno, Tulare, Kern, Kings and Madera Counties, in California, where it may be applied at 12 pints per acre for control of red scale by ground application. Do not use CPF 4E in combination with spray oil when temperatures are expected to exceed 95°F the day of application or for several consecutive days thereafter. Do not apply during the months of December, January or February.</p>			
Florida	aphids brown citrus aphid grasshoppers* orange dogs mealybugs scale insects snow scale Florida red scale purple scale long scale chaff scale black scale brown soft scale	2 - 8	ground: 100-1400 aerial: Min of 20
Restriction: Do not use a spray concentration of CPF 4E of less than 1/2 pt/ gal of water per acre.			
	citrus rust mites	4 - 8	100-700
Restriction: Do not use a spray concentration of CPF 4E of less than 1 pt/100 gal of water per acre.			
*Lubber grasshoppers must be controlled when they are small (less than 1 inch in length) by direct contact spray.			
Texas	aphids brown citrus aphid cutworms katydids mealybugs scale insects brown soft scale California red scale chaff scale	4 - 8	200-700
Texas	citrus rust mites (suppression)	4 - 8	200-700
Restriction: Do not use less than 1/2 pt of CPF 4E per 100 gallons of water in dilute applications.			
Crop: Small Transplanted Grapefruit, Orange and Other Citrus Trees			
Texas	aphids brown citrus aphid cutworms katydids mealybugs scale insects brown soft scale California red scale chaff scale	Max of 8	200-700
Remarks: Apply CPF 4E at a rate of 1 fl oz/1 gal of water with a backpack sprayer. Apply to runoff.			

CPF 4E may be tank mixed with ethion, dicofol, Agri-Mex, or Vendex. See "Mixing Directions" under the General Information Section. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for the other products used in combination with CPF 4E

Precautions: Observe local use directions for tank mix combinations especially in regard to applications of CPF 4E plus spray oil. Consult with a County Farm Advisor, County Agent, Extension Service Personnel, Agricultural Commissioner, or Pest Control Advisor for such information regarding a given locality.

Do not apply when trees are stressed by drought or high temperatures.

Do not tank mix CPF 4E with Difolatan 80, as crop injury may occur.

Restrictions: Do not apply more than 2 applications or more than 15 pints (7.5 lbs ai chlorpyrifos) of CPF 4E per acre per year (see additional application rate restrictions for specific California Counties). Do not make second foliar application within 30 days of the first application. Do not treat within 21 days of harvest for applications of up to 8 pints of Riot 4E per acre or within 35 days for application of rates above 8 pints per acre Do not do any work involving contact with trees within 4 days after treatment, Do not allow livestock to graze in treated areas.

Citrus Orchard Floors

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless Personal Protective Equipment (PPE) required for early entry is worn.

Ants (Excluding fire, harvester, carpenter, and pharaoh ants)

Use CPF 4E to control ants by applying the specified dose in 25 or more gallons of water with ground application equipment that will uniformly apply the spray to the orchard floor. To control foraging ants and suppress mounds, apply CPF 4E to the orchard floor at the rate of 3/4 to 1 quart per acre. Re-treat as needed. For best insect control, uniform coverage of the orchard floor is necessary. Do not apply where weed growths or other obstructions would impede uniform coverage of the orchard floor. Do not apply in tank mixtures with Evik herbicide. Do not allow spray to contact foliage. Foliar applications of CPF 4E or other products containing chlorpyrifos may be made in addition to the orchard floor treatments but must comply with the 10 day re-treatment interval.

CPF 4E may also be applied to citrus orchard floors through sprinkler irrigation systems only if the system uniformly covers the soil surface at the base of the tree. Use the specified amount of CPF 4E per acre. See "Sprinkler Irrigation" under the Product Information Section for further information.

Application With Dry Bulk Fertilizer: For impregnating CPF 4E on dry fertilizers, use a closed rotary drum mixer equipped with suitable spraying equipment. The spray nozzle should be positioned inside the mixer to provide uniform spray coverage of the tumbling fertilizer. Apply Plot 4E at the rate of 1 1/2 to 2 pints per acre to control ants in citrus orchard floors. The maximum concentration of CPF 4E to be added is 2 pints per 200 pounds of fertilizer. At the higher concentration of CPF 4E the fertilizer may not readily absorb all of the liquid. For a suitable free-floating mixture, an absorptive powder such as Micro-Cel E should be added separately and uniformly to the

fertilizer blend following addition of CPF 4E. Bulk fertilizers impregnated with CPF 4E should be applied immediately, not stored. All bulk containers should be tightly covered while the products are being transported and applied to reduce the chance of loss of CPF 4E via volatilization. Foliar applications of CPF 4E may be made in addition to the orchard floor treatments, but must comply with the 10 day re-treatment interval.

Compliance with any and all federal and state laws and regulations relating to the CPF 4E and fertilizer mixture is the responsibility of the person offering such mixture for sale or distribution.

Restrictions: Do not apply last treatment within 28 days before harvest. Do not apply more than 3 quarts of CPF 4E per acre per season. Do not make more than 3 applications. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not make second application of CPF 4E or any other product containing chlorpyrifos within 10 days of the previous application. Do not allow livestock to graze in treated areas.

Cranberries

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E by application as a broadcast, foliar spray to control brown spanworm, cranberry fruitworm, cranberry weevil, cutworms, fireworms, and Sparganothis fruitworms at the rate of 3 pints per acre. Mix the specified dosage in enough water to ensure thorough coverage and apply no less than 5 gallons of spray per acre when using ground equipment. For weevil control, apply once at flower bud development (late May, early June) and, if weevils are present, once after 100% bloom (early to mid July). For other insects, treat when field counts indicate damaging insect populations are developing or present. Apply only after the winter flood has been removed. To avoid pesticide contamination of floodwaters, make no applications while bogs are flooded.

CPF 4E may also be applied through sprinkler irrigation systems to control the above listed pests. For best results, use the specified rate of CPF 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation under the General Information Section for further information

Restrictions: Do not apply within 60 days before harvest. Do not make more than 2 applications per year for a total of 6 pints CPF 4E. Maximum single application rate is 3 pints CPF 4E (1.5 lbs ai chlorpyrifos) per acre. Do not make a second application of CPF 4E or any other product containing chlorpyrifos within 10 days of the first application.

Field Corn, Sweet Corn (Including Corn Grown For Seed)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless Personal Protective Equipment (PPE) required for early entry is worn.

For use to control cutworms, armyworms, corn earworm, corn rootworm adults, chinch bugs, grasshoppers, wireworms, flea beetle larvae and adults, aphids, billbugs, grubs, western bean cutworm, corn borers, symphylans, common stalk borer, and lesser cornstalk borer.

Preplant Incorporation Treatment

Use CPF 4E at the following rates by application in sufficient water to the soil surface and incorporate into the soil.

Pests	Dosage CPF 4E
cutworms symphylans	2 pt/acre

Use specified rate in not less than 10 gallons of water per acre and apply as a broadcast spray to the soil surface using suitable power operated ground spray equipment. On the same day of treatment, incorporate the insecticide into the top 2 to 4 inches of soil using a disc, field cultivator, or equivalent equipment.

CPF 4E may also be applied in tank mixtures with non-pressure fertilizer solutions and/or with Bladex®, Eradicane®, Sutan, Lasso®, Dual®, and atrazine herbicides. See "Mixing Directions" under the Product Information Section. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for the other products used in combination with CPF 4E

Preplant, At-Plant, or Preemergence Treatment In Conservation Tillage

Use CPF 4E at the following rate by application in sufficient water to surface trash and exposed soil:

Pests	Dosage CPF 4E
cutworms armyworms	1-2 pt/acre

Use specified rate in not less than 20 gallons of water per acre and apply as a broadcast spray using suitable power operated ground spray equipment. Use higher rates for residual control.

CPF 4E may also be applied in tank mixtures with non-pressure fertilizer solutions and/or with paraquat and Roundup herbicide, See "Mixing Directions" under the Product Information Section. Read and carefully follow all applicable directions, restrictions, and precautions, on labeling for the other products used in combination with CPF 4E.

T-Band At Plant Treatment

CPF 4E insecticide may be applied as a liquid T-Band in fields with no more than 30 percent cover of crop residue remaining on the soil surface. Apply CPF 4E as a liquid T-Band over an open seed furrow and incorporate into the top one-inch of soil using tines, chains or other suitable equipment. Position a flat fan nozzle behind the planter shoe, in front of the press wheel adjusted to provide a 5 to 6 inch bandwidth centered over the row. Apply CPF 4E at a rate of 2.4 fluid ounces per 1,000 linear feet of row (2 pints per acre with 40 inch row spacing) in a minimum spray volume of 5 gallons per acre.

If row spacing is less than 40 inches, the rate of CPF 4E must be adjusted so that no more than 2 pints per acre is applied. For example, for 30-inch rows, apply CPF 4E at a rate of 1.8 fluid ounces per 1000 linear feet of row (2 pints per acre).

Cultivation Time Treatment

Use CPF 4E at the rate of 2 pints per acre to control corn rootworm larvae. Apply CPF 4E as a water emulsion on both sides of the row at the base of the plants just ahead of the cultivator shovels. Cover the insecticide with soil around the brace roots. The best time to apply a basal treatment of a soil insecticide with cultivation is near the beginning of egg hatch. A cultivation application of CPF 4E may be made in addition to an at planting application of CPF 15G insecticide.

Postemergence Treatment

Use CPF 4E at the following rate by application in sufficient water to ensure thorough coverage of treated plants:

Pests	Dosage of CPF 4E
grasshoppers	1/2- 1 pt/acre
armyworms chinch bugs aphids corn rootworm adults Cutworms southern corn leaf beetle webworms western bean cutworm European corn borer (see note)	1 - 2 pt/acre

southwestern corn borer corn earworm	1 1/2 - 2 pt/acre
billbugs lesser cornstalk borer flea beetle adults common stalk borer	2 pt/acre

Note: The specified dosage will control silk clipping by corn rootworm adults. For European corn borer control, use 1 1/2 to 2 pints per acre when application is made with power-operated ground and aerial equipment and 1 to 2 pints per acre when application is made through a sprinkler irrigation system. See text below for generation specific treatment information.

Treat when field counts indicate that pests are or may become a problem. For best billbug, chinch bug, and flea beetle control, apply with sufficient water to ensure a minimum spray volume of 20 to 40 gallons per acre and 40 psi using ground spray equipment. On corn less than 6 inches tall, apply the insecticide spray in a 9 to 12 inch wide band over the row. On corn greater than 6 inches tall, apply the insecticide spray using drop nozzles directed to the base of the plant. Do not reduce the dosage for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone. When chinch bugs continue to immigrate to corn over a prolonged period or under extreme pressure, a second application of CPF 4E may be needed.

For cutworm, webworm, western bean cutworm, armyworm, aphid, European and southwestern corn borer, grasshopper, lesser cornstalk borer, corn rootworm adult, corn earworm, and common stalk borer control, apply as a broadcast spray using either aerial (fixed-wing or helicopter) or power operated ground spray equipment. For aerial application use 2 to 5 gallons of spray per acre. Control may be reduced at low spray volumes under high temperature and wind conditions. For cutworms, it is preferable to apply CPF 4E when soil is moist and worms are active on or near the soil surface. If ground is dry, cloddy, or crusty at time of treatment, worms may be protected from the spray and effectiveness will be reduced. If such conditions exist, shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment may improve control. Apply as needed to maintain control. Use higher rates for larger worms or when heavy cutworm infestations are expected or present. Fields should be monitored for cutworm presence or damage. A second application may be required if damage of density levels exceed economic thresholds established for your area. Consult your Agricultural Experiment Station or Extension Service Specialist for additional information concerning control practices in your area. For webworm control, shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment is necessary. For first-generation European corn borer control, treat when 25% to 50% of the corn plants show pinhole feeding or leaf-feeding scars. For maximum control potential, direct ground applications of CPF 4E into the corn leaf whorls. Scout fields within 5 days after application to determine if a second application is needed. University research indicates that achieving greater than 50% control of first-generation European borer with a single liquid insecticide treatment is highly dependent on timing, insecticide placement, and weather conditions. Treatment for control of second-generation European corn borer should be applied when field counts of egg masses indicate an infestation is present or about to develop. For southwestern corn borer control, treat when field counts of egg masses indicate pests are or may become a problem. A second application may be applied 10 to 14 days later, if needed due to reinfestation. For common stalk borer control, treat approximately 11 days after application of Roundup herbicide or after complete burn down with paraquat herbicide (3 to 5 days). Do not use CPF 4E in combination with the burn down herbicide for control of common stalk borer.

CPF 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar insects. Use the specified of CPF 4E in a tank mix with 2 pints per acre of non-emulsifiable oil. Maintain vigorous tank agitation to assure uniformity of the CPF 4E plus oil mixture throughout the injection period. CPF 4E may also be applied through sprinkler irrigation systems at the rate of 2 pints per acre to control corn rootworm larvae. Time application to coincide with the appearance of the second instar larvae. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. Apply with

enough water to wet the root zone to the depth control is needed. Under saturated soil conditions, allow enough soil drying to occur so that an application using a minimum water rate will not produce runoff. Consult university extension personnel or other experienced consultants to determine the need to treat and to aid in application timing. See "Sprinkler Irrigation" under the Product Information Section for further information.

Restrictions: Do not apply within 35 days before harvest of grain, ears, forage, or fodder. Do not apply more than 3 applications or a total of 6 pints of CPF 4E per acre per season, including the maximum allowed two granular applications at the 1 lb ai chlorpyrifos rate. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not make a second application of Lorsban-4E or other product containing chlorpyrifos within 10 days of the first application. If more than 1 lb ai granular chlorpyrifos per acre is applied at-plant (for a maximum of 1.3 lb ai per acre per season), only one additional application of a liquid product containing chlorpyrifos at 1 lb ai per acre is allowed per season, for a total of 2.3 lb ai chlorpyrifos per acre per season. Do not apply in tank mixes with Steadfast or Lightning herbicides. Do not aerially apply this product in Mississippi.

Sweet Corn Grown Only in Florida and Georgia

Use CPF 4E to control infestations of beet armyworm, fall armyworm, and corn earworm by application as a broadcast, foliar spray at the rate of 1 to 2 pints per acre. Mix the specified dosage in enough water to ensure thorough coverage and apply using suitable aerial or ground spray equipment. For aerial application, use at least 2 gallons of spray per acre. Treat when field counts indicate damaging pest populations are developing or present. Re-treat as necessary to maintain control but do not apply more than 3 applications per season.

CPF 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar insects. Use the specified rate of CPF 4E in a tank mix with 2 pints per acre of non-emulsifiable oil. Maintain vigorous tank agitation to assure uniformity of the CPF 4E plus oil mixture throughout the injection period. See "Sprinkler Irrigation" under the Product Information Section for further information.

Restrictions: Do not apply within 21 days before harvest of grain, ears, forage, or fodder. Do not apply more than 3 applications or a total of 6 pints of CPF 4E per acre per season, including the maximum allowed two granular applications at the 1 lb ai chlorpyrifos rate. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not make a second application of CPF 4E or other product containing chlorpyrifos within 10 days of the first application. If more than 1 lb ai granular chlorpyrifos per acre is applied at-plant (for a maximum of 1.3 lb ai per acre per season), only one additional application of a liquid product containing chlorpyrifos at 1 lb ai per acre is allowed per season, for a total of 2.3 lb ai chlorpyrifos per acre per season. Do not use in conjunction with post-plant broadcast, foliar applications of CPF 4E.

Cotton

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E for control of the following pests in all states except Arizona and California at the dosages indicated:

Pests	Dosage of CPF 4E
cotton fleahopper plant bugs (<i>Lygus. Mirids</i>)	3/8 - 1 pt/acre
fall armyworm grasshoppers thrips yellowstriped armyworm	1/2 - 1 pt/acre
cotton aphid	1/2 - 2 pt/acre

spider mites	1 pt/acre
beet armyworm cotton bollworm tobacco budworm cutworms pink bollworm salt marsh caterpillar	1 1/2 - 2 pt/acre

Use CPF 4E for control cotton pests in Arizona and California at the dosages indicated:

Cotton (California and Arizona)

Pests	Dosage of CPF 4E
armyworms cotton aphid cotton fleahopper <i>Lygus</i> salt marsh caterpillar silverleaf whitefly ¹ thrips	1 - 2 pt/acre
cotton bollworm cotton leaf perforator (suppression) tobacco budworm boll weevil cutworms pink bollworm spider mites (suppression)	2 pt/acre

¹ For control of silverleaf whitefly, apply in tank mix combination with the specified rate of a pyrethroid insecticide labeled for control or suppression of whitefly. Do not apply more than 3 applications per season.

Mix the required dosage with sufficient water to ensure thorough coverage of plants and apply using aerial or power operated ground spray equipment. For aerial application, use at least 1 gallon of spray per acre. Treat when field counts indicate damaging insect populations are developing or present. Re-treat as necessary to maintain control but do not apply more than 3 applications per season.

CPF 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. Use the specified rate of CPF 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" under the Product Information Section for further information.

For effective control of spider mites when large numbers of eggs are present, apply a second spray 10 days after initial treatment to control newly hatched nymphs or after 3-5 days with a non-chlorpyrifos products..

For best results on bollworms and budworms, it is suggested that fields be scouted twice per week and treatments made when worms are 1/4 inch or less in length. The following table illustrates the size of worms in relation to age and stage of development (instar) as a guide to timing of treatments for best control.

From the table it can be seen that a scouting schedule of only once per week will not be satisfactory since the worms may be too big to control effectively by the seventh or eighth day.

Timing for the Best Worm Control

	Age (Days)	Average Size	Instar
Get the worms at this stage	Hatch	1/16"	Hatch
	3	3/32"	I
	5	9/32"	II
	6	7/16"	III
	8	11/16"	IV

1/16" = --

3/32" = ---

9/32" = ----

(actual size pictures of worms to be inserted)

7/16" = ----

11/16" = ----- (approximately the size of a dime)

Proper application techniques help to ensure thorough spray coverage and correct dosage and are thus important in obtaining good control of pests. Consider these suggestions when applying CPF 4E on cotton.

Aerial Application

Shorten boom length to avoid spray entering the vortices at the wing tips. Swath width should be reduced when wind direction is the same as direction of spraying.

The proper nozzle arrangement and swath width to avoid skips and vortices effect can be checked out by flying over a paper tape (adding machine paper) using water with or without soluble dye.

(The dye gives a permanent record).

Flying at a height of 5 to 15 feet above the target results in the best coverage.

Nozzle orientation of the boom is important. More break-up occurs when nozzles are pointed straight down versus the straight back position. Desired droplet size (100 to 200 microns) can be obtained by angling the nozzles somewhere in this range.

Marking of swath by flagging or permanent markers is essential.

Ground Application

Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom; drift spray is wasted spray so do not depend on it. Use flat fan or disc-core hollow cone nozzles with maximum spacing of 20 inches and a spray pressure of 40 to 60 psi with a droplet size of 100 to 200 microns.

Restrictions: Do not apply within 14 days before harvest. Do not make more than three applications of CPF 4E or other product containing chlorpyrifos per crop season. Do not apply more than 6 pints of CPF 4E (3 lb ai chlorpyrifos) per acre per season. Maximum single application rate is 1 lb ai chlorpyrifos (2 pints of CPF 4E) per acre. Do not make a second application of Lorsban-4E or other product containing chlorpyrifos within 10 days of the first application. Do not allow meat or dairy animals to graze in treated areas. Do not feed gin trash or treated forage to meat or dairy animals.

Figs

(Only for use in California)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E at the rate of 2 quarts per acre for control of dried fruit beetle by application in sufficient water to the soil surface followed by incorporation into the top 3 inches of soil. Apply to fig orchard soil as a dormant application in late winter prior to beetle emergence and prior to leaf formation.

Restrictions: Make only 1 application per year. Maximum single application rate is 2 quarts (2 lbs ai chlorpyrifos) per acre. Do not apply within 210 days (7 months) of harvest. Based on available residue data, use of CPF 4E on figs is restricted to California.

Grapes

(Not for use in Mississippi. For use only in states east of the Rocky Mountains)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E for control of grape root borer by application just before the pest emerges from the soil. Mix 4 1/2 pints (2.25 lb a.i./A) of CPF 4E with 100 gallons of water and apply 2 quarts of the diluted spray mixture to the soil surface on a 15-square foot area around the base of each vine. Do not allow spray to contact fruit or foliage.

Restrictions: Do not apply within 35 days before harvest. Do not make more than 1 application per season. Maximum single application rate is 4.5 pints CPF 4E (2.25 lb ai chlorpyrifos) per acre. Based upon available residue data, the use of CPF 4E in grapes is restricted to states east of the Rocky Mountains.

Mint (Peppermint)

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E by application as a broadcast, foliar spray to control cutworms at the rate of 2 to 4 pints per acre and mint root borer at the rate of 4 pints per acre. Mix the specified dosage in water to give no less than 10 gallons of spray per acre and apply using ground spray equipment. For cutworm control, treat during May and June when field counts indicate damaging insect populations are developing or present. When larvae are less than 3/4 inch in length, use the 2 pint rate. When larvae are 3/4 inch or more in length, use the higher rate. Make only 1 application during the growing season. For mint root borer control, apply postharvest when field counts indicate damaging insect populations are developing or present. Follow treatment with approximately 1 acre-inch of sprinkler irrigation immediately after application to incorporate the insecticide into the soil. Make only 1 postharvest application per season.

CPF 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed pests. Use the specified rate of CPF 4E per acre. Maintain vigorous tank agitation throughout the injection period. See "Sprinkler Irrigation" under the Product Information Section for further information.

Restrictions: Do not apply within 90 days before harvest. Make only 1 application per year. Maximum single application rate is 4 pints CPF 4E (2 lbs ai chlorpyrifos) per acre.

Onions (Dry Bulb)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E to control onion maggot by application as an in-furrow drench. Apply CPF 4E at the rate of 1.1 fluid ounce(0.035 lb a.i./A) per 1,000 linear feet of row at 18-inch row spacing. Use a minimum of 40 gallons of total drench per acre. Incorporate to a depth of 1 to 2 inches.

Restrictions: Do not apply within 60 days of harvest. Do not make more than 1 application per year. Maximum single application rate is 0.03 lb ai chlorpyrifos per 1000 ft of row. Do not aerially apply this product in Mississippi.

Peanuts

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

For suppression of wireworms, apply CPF 4E at a rate of 4 pints per acre as a preplant broadcast spray to the soil surface followed by immediate soil incorporation to a depth of 3 to 4 inches. Use a minimum of 10 gallons of total spray per acre.

Restrictions: Do not apply within 21 days of harvest. Maximum single application rate is 4 pints CPF 4E (2 lb ai chlorpyrifos) per acre. The combined total of preplant and post-plant applications of CPF 4E and CPF 15G or any other products containing chlorpyrifos must not exceed 4 pounds active ingredient per acre per season. Do not make more than 1 application per season. Do not feed treated peanut forage or hay to meat or dairy animals. Do not apply aerially.

Sorghum - Grain Sorghum (Milo)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E insecticide for control of the following pests at the Dosages indicated:

Pests	Dosage of CPF 4E	Specific Directions
sorghum midge	1/2 pt/acre	Apply when 30% to 50% of the seed heads are in bloom. Repeat at 3-day intervals if necessary.
grasshoppers yellow sugar cane aphid and other aphids	1/2 -1 pt/acre	

greenbug	1/2 - 2 pt/acre	For infestations of greenbug that are difficult to control, use a higher dose within the indicated dose range.
chinch bugs lesser cornstalk borer	1 - 2 pt/acre	Apply as a directed spray toward the base of the plant using power operated ground spray equipment with sufficient water to ensure coverage of an 8-12 inch band centered in the row. On plants less than 6 inches high, apply a 8-12 inch band over the row. Do not reduce the dosage for banded or directed applications. Concentrate the full- labeled dosage rate in the treated zone.
webworms	1 pt/acre	
armyworms cutworms	1 - 2 pt/acre	
European and southwestern corn borer	1 1/2-2 pt/acre	
corn earworm	2pt/acre	

Mix the specified dosage in enough water to ensure thorough coverage and apply using suitable aerial or ground spray equipment.

To minimize chemical injury, do not apply CPF 4E to drought stressed grain sorghum within 3 days following irrigation or rain except where the product is applied in irrigation water.

CPF 4E may also be applied through sprinkler irrigation systems as a post emergence broadcast application to control the above listed foliar pests. For best results, use the recommended rate of CPF 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" under the General Information Section of this label.

Precaution: Be aware that sorghum lines used in seed production fields may be more sensitive to chemical injury. Susceptible inbred lines or hybrids are likely to be at greater risk of yield-reducing chemical injury when sprayed at the higher rates of application. Do not apply more than 1 pint per acre of CPF 4E to seed sorghum if the additional risk of crop injury is unacceptable.

Restrictions: The treated crop must not be used for grain, forage, fodder, hay or silage within 30 days after application of 1 pint of CPF 4E per acre or within 60 days after application of rates greater than 1 pint per acre. Do not treat sweet varieties of sorghum. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not apply more than 3 applications or a total of 3 pints (1.5 lb ai chlorpyrifos) of CPF 4E or any other product containing chlorpyrifos per acre per season. Do not make second or third treatment within 10 days of a previous treatment. Do not aerially apply this product in Mississippi.

Soybeans

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

For use to control armyworms, bean leaf beetle, corn earworm, cutworms, European coat borer, grasshoppers, green cloverworm, lesser cornstalk borer. Mexican bean beetle, saltmarsh caterpillar and other woollybears, southern green stink bug, spider mites, and velvetbean caterpillar.

Soil Treatment

Use CPF 4E at the rate of 1 to 2 pints per acre to control cutworms and lesser cornstalk borer. Mix the specified dosage in a minimum of 10 gallons of spray per acre and apply to the soil surface using suitable ground spray equipment. Equivalent rates of insecticide spray required per 100 feet of row for various row spacing are given in the accompanying table. For at-plant treatments apply the insecticide over the row in a 4 to 6 inch band in front of the planter shoe or press wheel or after the press wheel followed by a drag chain for light incorporation. Do not apply as an in furrow treatment. For postemergence rescue treatments, apply as a directed spray in a 9 to 12 inch band at the base of the plant. To plants less than 6 inches high apply over the top in a 6 to 12 inch band. Treat when field counts or conditions indicate that pests are or may become a problem.

	Fluid Ounces of Spray Required Per 100 Feet of Row for various Row Spacings			
Volume of Spray per Acre	36"	32"	28"	24"
10 gallons	8.8	7.9	6.9	5.9
15 gallons	13.2	11.8	10.3	8.8
20 gallons	17.6	15.7	13.7	11.8

Foliar Treatment

Use CPF 4E at the following rate by application in sufficient water to ensure thorough coverage of treated plants:

Pests	Dosage of CPF 4E
European corn borer southern green stink bug	2 pt/acre
bean leaf beetle cutworms corn earworm saltmarsh caterpillar and other woolly bears	1-2 pt/acre
Mexican bean beetle armyworms	1 – 1 1/2 pt/acre
velvetbean caterpillar grasshoppers green cloverworm spider mites	1/2 - 1 pt/acre

Apply as a broadcast spray using either aerial or ground equipment when field counts indicate damaging insect populations are developing or present; re-treat as necessary to maintain control. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days after initial treatment to control newly hatched nymphs. On determinate soybeans do not apply more than 1 application after pod set.

CPF 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. Use the specified rate of CPF 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See "Sprinkler Irrigation" section under the Product Information Section of this label for further information.

Restrictions: Do not apply more than 6 pints of CPF 4E per acre or 3 pounds of chlorpyrifos (active ingredient) per acre per season, or make more than 3 applications per season. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not apply within 28 days before harvest. Do not make a second or third application within 14 days of a previous application. Do not allow livestock to graze in treated areas or otherwise feed treated soybean forage, hay, and straw to meat or dairy animals.

Strawberries

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E by application as a broadcast foliar spray to control strawberry bud weevil at the rate of 1 quart per acre. Apply in a minimum of 40 gallons of spray per acre when buds first appear and 10 to 14 days later. Do not apply after berries start to form or when berries are present. CPF 4E should not be tank mixed with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination non-injurious under your current conditions of use. Phytotoxicity may occur when CPF 4E is applied to strawberries experiencing high temperature and drought stress.

Restrictions: For pre-bloom use only. Maximum single application rate is 1 quart CPF 4E (1 lb ai chlorpyrifos) per acre. Do not make more than 2 applications per season or apply within 21 days before harvest.

Sugar Beets

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Soil Treatment (At Planting or Preplant Incorporated)

To reduce feeding damage from early season insects such as cutworms, use CPF 4E at planting or as a preplant treatment and incorporate to a depth of 1 to 2 inches. Do not apply as an in-furrow treatment. Apply 1 pint of CPF 4E per planted acre to a 10-inch wide band centered on the row for furrows 30 inches apart. (For rows 30 inches apart, this is equivalent to 9.2 fluid ounces of CPF 4E per 10,000 feet of row). For other row widths, adjust the spray volume per planted acre in proportion to the area actually treated.

Postemergence Treatment

Apply CPF 4E as a broadcast or banded foliar spray. Treat when field counts indicate that damaging insect populations are developing or present. Do not make more than 3 applications per season.

Broadcast Application: Apply the specified dosage in water using 2 to 5 gallons of finished spray per acre when using aerial spray equipment or 10 to 30 gallons per acre when using ground spray equipment.

Broadcast Application Through Sprinkler Irrigation Systems

CPF 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the foliar pests. Use the specified rate of CPF 4E per acre. Maintain vigorous tank agitation to assume uniformity of the application through the injection period. See "Sprinkler Irrigation" under the Product Information Section for further information.

Band Application: Apply the specified dosage within the band using a minimum of 6 1/2 gallons of finished spray per acre. Apply the spray in a 5 to 7 inch wide band over the row. Do not reduce the dosage for band applications. Concentrate the full-labeled dosage rate in the treated zone. For best results, band-applied treatments should be lightly incorporated either mechanically or with irrigation.

Use CPF 4E at the rates indicated to control the listed pests.

Dosage of CPF 4E			
Pests	Broadcast	Band	Timing/Special Directions
grasshoppers	1/2 - 1 pt/acre		Low rate will control small nymphs (1st through 3 rd instar).
spider mites	1 pt/acre	2/3 pt/acre	
fall armyworm yellowstriped armyworm webworms	1 - 2 pt/acre	2/3 - 1 1/3 pt/acre	
beet armyworm	1 1/2 – 2 pt/acre	1- 1 1/3 pt/acre	
cutworms flea beetle adults	2 pt/acre	1 1/3 pt/acre	
Sugar beet maggots adult ¹	1/2 - 1 pt/acre	-	To target adults present at the time of application based on local field trap monitoring, apply anytime from 7 days before until 3 days after peak adult emergence.
sugar beet root maggot larvae ¹	2 pt/acre	2/3 – 1 1/3 pt/acre	Use as supplemental treatment following an at-plant insecticide treatment for control of root maggot. Application timing should be based on local field trap monitoring. Apply anytime from 7 days before until 3 days after peak adult emergence.
Suger beet root maggot larvae ¹	-	1 1/3 – 2 pt/acre	Use as primary treatment to control root maggot. Application timing should be based on local field trap monitoring. Apply anytime from 7 days before until 3 days after peak adult emergence.

¹ To prevent potential development of insecticide resistance in sugar beet root maggot, producers are encouraged to take the following steps: 1) do not apply more than 2 applications of CPF 4E per season when adults are active; 2) if an organophosphate insecticide was applied at planting, make no more than 1 postemergence application of CPF 4E when adults are active.

Restrictions: Do not apply within 30 days of harvest of beet roots and tops. Do not apply more than a total of 6 pints per acre of CPF 4E on a broadcast basis per season, or make more than 3 applications per season. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not make a second or third application within 10 days of a previous application. Do not allow livestock to graze in treated areas or harvest treated beet tops as feed for meat or dairy animals within 30 days of last treatment.

Sunflowers

(Not for use in Mississippi)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

For use to control cutworms, sunflower beetle larvae and adults, stem weevil, sunflower moth, woolly bears, seed weevil, and grasshoppers.

Preplant Incorporation Treatment

Use CPF 4E at the following rates by application in sufficient water to the soil surface and incorporate into the soil:

Pests	Dosage of CPF 4E
Cutworms	2 – 4 pt/acre

Use recommended rate in not less than 10 gallons of water per acre and apply as a broadcast spray to the soil surface using suitable power operated ground spray equipment. On the same day of treatment, incorporate the insecticide into the top 2 to 4 inches of soil using a disc, field cultivator, or equivalent equipment.

Restrictions: Do not apply more than one application per season. Do not harvest within 42 days of application. Maximum single application rate is 4 pints CPF 4E (2 lb ai chlorpyrifos) per acre. Grazing of meat or dairy animals in treated orchards is prohibited.

Postemergence Treatment

Use CPF 4E for control of the following pests at the dosage indicated by application in sufficient water to ensure thorough coverage of treated plants:

Pests	Dosage of CPF 4E
cutworms sunflower beetle larvae and adults stem weevil sunflower moth banded sunflower moth woolly bears seed weevil	1 1/2 pt/acre
grasshoppers	1 pt/acre
Tarnished plant (Lygus) bugs	1 -1 1/2 pt/acre

Apply as a broadcast spray using either aerial (fixed-wing or helicopter) or power-operated ground spray equipment when field counts indicate that pests are or may become a problem. For cutworm control, a second treatment may be made 7 to 10 days later, if needed. For stem weevil control, optimal treatment time is within 5

to 7 days after adult weevils begin to appear. For sunflower moth control, make first application during early 1% to 5% bloom stage. A second treatment may be made 7 days later, if needed. For seed weevil control, treat when field counts indicate there are 10 to 12 adults per plant for oil crops and 1 to 3 adults per plant on confectionery crops. Additional treatments should be made at successive 7 to 10 day intervals if field counts indicate need to retreat. For sunflower beetle larvae or adult control, treat when field counts indicate there are 10 larvae or 1 to 2 adults per seedling. Additional treatments may be made at successive 7-to-10-day intervals if field counts indicate need to re-treat.

Restrictions: Do not apply more than 3 applications or a total of 6 pints of CPF 4E per acre per season. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre per season. Do not apply within 42 days before harvest. Do not allow livestock to graze in treated areas. If a preplant application has been made at the 4 pint rate, then only one postemergence treatment is allowed.

Sweet Potatoes

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E to reduce the feeding damage caused by populations of *Conderus* wireworm, *Systema* flea beetle, and the sweet potato flea beetle. Apply at the rate of 4 pints per acre as a broadcast (overall) spray to the soil surface followed by incorporation. Mix the specified dosage with enough water to obtain uniform coverage and apply as a coarse spray using suitable ground spray equipment. Incorporate the insecticide to a depth of 4 to 6 inches as soon as possible after application by using a rotary hoe, disc cultivator, or other suitable incorporation equipment. Plant the crop in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time that CPF 4E will protect against feeding damage). CPF 4E will not control false wireworms or white fringe beetle or other grubs that attack sweet potatoes.

Restrictions: Do not make more than 1 application per season. Maximum single application rate is 4 pints CPF 4E (2 lb ai chlorpyrifos) per acre. Do not harvest within 125 days of treatment. Do not apply aerially in Mississippi.

Tobacco

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E as a preplant treatment to control larvae of cutworms, flea beetles, mole crickets, root maggots, and wireworms. Apply 2 pints of CPF 4E per acre in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate the insecticide into the soil to a depth of 2 to 4 inches using suitable equipment. The application of CPF 4E will also suppress the movement of imported fire ants into treated fields.

To control the above insects and low to moderate populations of rootknot nematodes in North Carolina, South Carolina, and Virginia, use CPF 4E at the rate of 2 pints per acre. To control the above insects and moderate populations of rootknot nematodes in all tobacco growing regions, use CPF 4E in a tank mix with Nematicur 3 at the rate of 2 pints of CPF 4E plus 8 pints of Nematicur 3 nematicide per acre. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for Nematicur 3 used in combination with CPF 4E. Apply the specified dosage in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil to a depth of at least 4 inches using suitable equipment. Where the nematode species *Meloidogyne arenaria* or *M.javanica* are present or high populations of *M.incognita*, apply Telone II soil fumigant at the specified label rate.

Before broadcast application of CPF 4E on to existing beds, knock down beds to final shape for transplanting. Use of PTO-driven implements that will incorporate CPF 4E to a depth of 4 inches is recommended.

Restrictions: Do not make more than 1 application per season. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not apply aerially in Mississippi.

Tree Fruits

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless Personal Protective Equipment (PPE) required for early entry is worn.

Apples, pears, plums, prunes: Use CPF 4 E as a dormant or delayed dormant spray at the rates indicated to control the following insects on the crops listed. While Plot 4E may be used without oil, oil is recommended to control additional pests such as European red mite.

For dilute sprays, tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using suitable ground spray equipment. [See "Additional Precautions Specific to California" (below) for use in California].

For low volume (concentrate) sprays, less than 200 gallons of spray mixture per acre use the same amount of CPF 4E as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of CPF 4E for severe infestations. Use oil as recommended by your State Agricultural Experiment Station or Extension Service Specialist.

Precautions

Because cold or dry conditions may cause CPF 4E plus oil sprays to infuse trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated. Do not use more than 4 pints of CPF 4E per acre.

Additional Precautions Specific to California: Use a minimum of 250 gallons of total spray volume per acre. Do not use more than 4 gallons of spray oil per acre on almonds, peaches, or nectarines. Do not use any adjuvants or surfactants in addition to or as a substitute for petroleum spray oil in a tank mix with CPF 4E. Do not apply on almonds in the following counties in California: Butte, Colusa, Glenn, Solano, Sutter, Tehama, Yuba, and Yuba.

Restrictions: Post-bloom applications to apple trees are prohibited, Make only 1 application during the dormant season. Maximum single application rate is 1 pint CPF 4E (0.5 lb ai chlorpyrifos) per 100 gallons of water. Do not allow meat or dairy animals to graze in treated orchards.

Cherries: Use CPF 4E for the control of lesser peach tree borer, and American plum borer by application as a trunk spray. Mix 1 1/2 to 3 quarts of CPF 4E with 100 gallons of water and apply as a course, low pressure spray to give uniform coverage of tree trunks and lower limbs. Make a second application 2 weeks after the first one and a third application after harvest. Avoid contact with foliage in sweet cherries as premature leaf drop may result. Consult your State Agricultural Experiment Station or Extension Service Specialist for proper time to treat in your area.

In addition, 1 of the 3 allowable applications per year may be applied as a dormant spray of San Jose scale, peach twig borer, and climbing cutworms. For control of these pests, tank mix 1/2 to 1 pint of CPF 4E with 1 to 2 gallons of a petroleum oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using ground spray equipment. For low volume (concentrate) sprays (40 to 100 gallons of spray mixture per acre) use the same amounts of CPF 4E and spray oil per acre required for application as a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of CPF 4E for severe infestations. Use oil as recommended by your State Agricultural Experiment Station or Extension Service Specialist.

Restrictions: Make only 3 applications per year. Do not apply within 6 days before harvest. Do not allow meat or dairy animals to graze in treated orchards.

Nectarines, Peaches: Use CPF 4E for the control of peach tree borers by application as a trunk spray before newly hatched borers enter the trees. Mix 3 quarts of CPF 4E with 100 gallons of water and apply as a course, low-pressure spray to give uniform coverage of tree trunks. Thoroughly wet all bark areas from ground level to scaffold limbs. Do not allow spray to contact fruit. Consult your State Agricultural Experiment Station's or Extension Service Specialist's written recommendations for proper time to treat in your area.

CPF 4E may also be used as a preplant dip application for nonbearing peach trees at the equivalent application rate of 3 quarts per 100 gallons of water for control of peach tree borer. Dip trees several inches above the grafting bud scar and plant immediately or allow drying before returning to storage. Do not allow peach trees to remain in contact with the dip solution.

Restrictions: Make only 1 application per season. Do not apply within 14 days before harvest. Do not allow meat or dairy animals to graze in treated orchards.

Use CPF 4E at the rates indicated to control the listed pests.

Crop/Pest	CPF 4E per 100 Gallons of Spray ¹	Application Restrictions
Apples rosy apple aphid San Jose scale <i>Lygus</i> pandemis leafroller climbing cutworms oblique banded leafroller	1 pt/100 gal (1200-600 gal Finished spray/A) to 1 to 3 lb a.i./A	Spray application Dormant/Delayed dormant. Ground Equipment. Do not apply more than 1 application per season. Application may be made alone or as a tank mix with petroleum spray oil. Grazing of meat or dairy animals in treated orchards is prohibited.
Cherries (sweet) Lesser peach borer American plum borer San Jose scale Peach twig borer climbing cutworms	3 qt./100 gal	Trunk spray (bark) application. Foliar and postharvest and/or dormant/delayed dormant. Ground Equipment. Do not apply more than 3 applications (9 lb a.i./A) per season. One of the three applications per season may be applied as a dormant spray tank mixed with petroleum spray oil at 0.5 lb a.i./100 gal. Do not harvest within 6 days of application. Grazing of livestock in treated orchards is prohibited.
Pears San Jose scale climbing cutworms pear psylla adults	1 pt/100 gal (200-600 gal finished spray/A) 1 to 3 lb a.i./A	Spray application Dormant/Delayed dormant. Ground Equipment. Do not apply more than 1 application per season. Application may be made alone or as a tank mix with petroleum spray oil. Grazing of meat or dairy animals in treated orchards is prohibited.

Nectarines Peaches Plum/Prunes San Jose scale mealy plum aphid climbing cutworms peach twig borer	1 pt/100 gal (200-600 gal finished spray/A) 1 to 3 lb a.i./A	Spray application Dormant/Delayed dormant. Ground Equipment. Do not apply more than 1 application per season. Application may be made alone or as a tank mix with petroleum spray oil. Grazing of meat or dairy animals in treated orchards is prohibited.
--	---	---

¹ Based on 200 to 600 gallons per acre as a dilute spray.

Tree Nut Crops

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E at the dosages indicated by application as a foliar spray to control pests listed in the following table. Mix the required dosage in sufficient water to ensure thorough and complete coverage of the foliage and crop and apply as a concentrate or dilute spray using conventional, power-operated spray equipment. For dilute sprays applied to tree nut crops, mix the required dosage in sufficient water to allow for spray to runoff. For concentrate sprays, apply an equivalent amount of CPF 4E per acre. Treat when pests appear or in accordance with local conditions. Insect control by aerial application may be less than control by ground application because of less coverage. Consult your State Agricultural Experiment Station, certified Pest Control Advisor, or Extension Service Specialist for specific use information in your area.

Almonds, Filberts, Walnuts

Use CPF 4E at the rates indicated to control the listed pests.

Crop	Insects Controlled	Dosage CPF 4E
almonds	navel orangeworm peach twig borer San Jose scale	4 pt/acre
filberts	eye-spotted bud moth filbert aphid filbert leafroller filbert worm oblique-banded leafroller omnivorous leaf-tier winter moth	3 - 4 pt/acre
walnuts	codling moth walnut husk fly walnut scale	4 pt/acre
Restrictions: Do not apply more than 2 foliar applications or a total of 8 pints of CPF 4E per acre per season on almonds, filberts, and walnuts. Maximum single application rate is 4 pints CPF 4E (2 lbs ai chlorpyrifos) per acre. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated orchards.		

Pecans

Use CPF 4E at the rates indicated to control the listed pests.

Insects Controlled	Dosage of CPF 4E (Dilute or Concentrate)	Remarks
spittlebugs	1 - 4 pt/acre	Use a dosage of 2 to 4 pints per acre for concentrate sprays.
pecan nut casebearer fall webworm	1 1/2 - 4 pt/acre	
<i>Phylloxera</i> spp. ¹ black pecan aphid hickory shuckworm ² pecan leaf scorch mite (suppression) ³ fire ants and other ant species ⁴	2 - 4 pt/acre	¹ For best <i>Phylloxera</i> s spp. control, make 2 applications at a 7 to 10 day interval using a minimum of 1.0 pint of CPF 4E per acre starting at bud swell. ² For best results make 2 applications 10 to 14 days apart. ³ To suppress pecan leaf scorch mite, use a preventative program. ⁴ For ant control, apply as an orchard floor spray. Do not apply where weed growth or other obstructions prevent uniform coverage of the orchard floor.
yellow pecan aphid black margined aphid	1 -4 pints of CPF 4E plus: 5.33 fl. oz. of Pydrin 2.4E, or 1 .7 0 fl. oz. of Asana 1.9EC, or 3.00 fl. oz. of Ammo 2.5EC, or 2.56 fl. oz. of Cymbush 3E	

Restrictions: Do not apply more than 5 applications or a total of 8 pints of CPF 4E per acre per season on pecans. Maximum single application rate is 4 pints CPF 4E (2 lbs ai chlorpyrifos) per acre. Do not apply within 28 days of harvest. Do not allow livestock to graze in treated orchards. Make no applications of tank mixtures closer to harvest than the longest preharvest interval shown for any of the products in the tank mixture. For dilute applications with ground equipment use at least the minimum rate of CPF 4E listed for the pest. Apply in 100 to 600 gallons of water per acre. For aerial applications use 5 to 15 gallons of water per acre. Note: With aerial application, control may be reduced due to poor coverage. Up to 10 pints of CPF 4E may be applied per acre per year.

Almond and Walnut Orchard Floors

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E to control ants (excluding fire, harvester, carpenter, and pharaoh ants) by applying the specified dose with ground application equipment that will uniformly apply the spray to the orchard floor. Use when ant activity becomes evident within the orchard. Since worker ants cease most of their foraging activity at temperatures above 90°F, best results will be achieved with applications made at temperatures below 90°F at

the time of application. CPF 4E may also be applied to almond and walnut orchard floors through sprinkler irrigation systems only if the system uniformly covers the soil surface. Dosage of CPF 4E and spray volume may vary depending on the irrigation method employed in the orchard as follows:

Ant Control in Sprinkler or Drip Irrigated Orchards

Apply CPF 4E as a broadcast spray to the entire orchard floor using ground spray equipment at 4 to 8 pints per acre in 25 or more gallons of water. Use the high rate for heavy infestations and the low rate for light infestations. In orchards where ant activity is concentrated around the irrigation emitters, apply the high rate to a 6 to 8 foot band along the drip irrigation line and the low rate to the rest of the orchard.

Ant Control in Flood Irrigated Orchards

Apply CPF 4E at 4 to 8 pints per acre in 25 or more gallons of water to the entire orchard floor using ground spray equipment. Apply the high rate to heavily infested areas and the low rate to lightly infested areas. Where ant colonies are abundant only in the berm areas, apply CPF 4E at 8 pints per treated acre in 50 or more gallons of water to a 6 to 10 foot band along the tree line (berm).

Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor, Mow or chemically control weeds before the application of CPF 4E. Foliar applications of CPF 4E may be made in addition to the orchard floor treatment.

Restrictions: Do not apply more than 2 applications or a total of 8 pints of CPF 4E per acre per season to the orchard floor. Maximum single application rate is 8 pints CPF 4E (4 lb ai chlorpyrifos) per acre. If the 8 pint rate is used, a second application is not allowed. Do not make a second application within 10 days of the first application. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated orchards.

Vegetables (*Brassica* (Cole) Leafy Vegetables¹ and Radish, Rutabaga, and Turnip)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 3 days for cauliflower and 24 hours for the other listed vegetables unless Personal Protective Equipment (PPE) required for early entry is worn.

Use CPF 4E at the dosages indicated to control the pests listed in the following table. To avoid phytotoxicity in vegetables, except Brussels sprouts, do not mix with other pesticide products or treat plants that are under extreme heat and drought stress.

Corn	Insects Controlled	Dosage of CPF 4E	Use Directions
broccoli cauliflower cabbage Chinese cabbage (bok-choy, napa)	root maggot root aphid	1.2 fl oz/ 1,000 linear ft of row	Soil band treatment at planting or Transplanting -or- Directed spray application Post-transplant. Ground Spray Equipment. For direct seeded crops apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time. Shallow incorporation is necessary. Placement behind the planter shoe and in front of the press wheel is recommended. For transplanted crops, apply CPF 4E as a water-based spray directed to the base of the plants immediately after setting. Use a minimum of 40 gallons of total spray per acre. Do not add any additional adjuvant, surfactants or spreader stickers. Do not apply as a foliage application.

Restrictions: Do not apply within 30 days of harvest. Do not apply more than 4 pints CPF 4E (2 lb ai chlorpyrifos) per acre for cauliflower, or 4.5 pints CPF 4E (2.25 lb ai chlorpyrifos) per acre for broccoli, cabbage, and Chinese cabbage (72 fl oz/A) per season. Do not make more than 1 application per season.

broccoli cabbage	root maggot root aphid	1.3 fl oz/1,000 linear ft of row	Soil injected side dress application. Apply CPF 4E in a water emulsion or with liquid fertilizer injected as a side dress on each side of the row after plants are established. Avoid mechanical damage to crop roots. Use a minimum of 15 gallons of total spray volume per acre.
---------------------	---------------------------	-------------------------------------	---

Restrictions: Do not apply within 30 days of harvest. Do not apply more than 4.5 pints CPF 4E (2.25 lb ai chlorpyrifos) per acre (72 fl oz/A) or make more than 1 application per season.

brussels sprouts collards kohlrabi turnips	root maggot root aphid	1.4 fl oz/ 1,000 linear ft of row	Soil band treatment at planting or Transplanting Directed spray application Post-transplant. Ground Spray Equipment. For direct seeded crops apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time. Shallow incorporation is necessary. Placement behind the planter shoe and in front of the press wheel is recommended. For transplanted crops, apply CPF 4E as a water-based spray directed to the base of the plants immediately after setting. Use a minimum of 40 gallons of total spray per acre. Do not add any additional adjuvant, surfactants or spreader stickers. Do not apply as a foliage application
---	---------------------------	---	--

Restrictions: Do not apply within 30 days of harvest. Do not apply more than 4.5 pints CPF 4E (2.25 lb ai chlorpyrifos) per acre (72 fl oz/A) or make more than 1 application per season.

brussels sprouts	Armyworms Cabbage aphid Cutworms Imported Cabbage-worm Striped flea Beetle (adult)	1 lb/A	Broadcast application foliar. Ground or Aerial Equipment. Apply in 20 to 150 gallons of water per acre. Apply when insects appear on foliage and at 7 to 14 day intervals thereafter as needed. Consult your State Agricultural Experiment Station, Extension Service Specialist or Integrated Pest Control Advisor for proper time to treat your area.
Restrictions: Do not apply within 21 days of harvest. Do not make more than 3 applications per year. Maximum single application rate is 2 pints CPF 4E (1 lb ai chlorpyrifos) per acre. Do not apply more than 3 lb a.i./A (96 fl oz/A) (max. 3 applications) per season.			
radishes	root maggot	0.5 fl oz/1,000 linear ft of row	Apply the specified dosage as a water-based drench in the seed furrows, with the seed at planting time. Use a minimum of 40 gallons of total drench per acre.
Restrictions: Do not apply within 30 days of harvest. Do not apply more than 5.5 pints CPF 4E (2.75 lb ai chlorpyrifos) per acre (96 fl oz/A) or make more than 1 application per season.			
rutabagas	root maggot	1.6 fl oz/1,000 linear ft of row	Apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time, behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Use a minimum of 40
Restrictions: Do not apply within 30 days of harvest. Do not apply more than 4.5 pints CPF 4E (2.25 lb ai chlorpyrifos) per acre (72 fl oz/A) or make more than 1 application per season. The use of rutabaga tops for food/feed purposes is prohibited.			

Wheat

(For use only in Arizona, California, Colorado, Idaho, Kansas, Minnesota, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington and Wyoming).

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless Personal Protective Equipment (PPE) required for early entry is worn.

For use to control aphids (including Russian wheat aphid), wheat midge, brown wheat mite, grasshoppers, army cutworms and to provide suppression of other cutworm species.

Pests	Pilot4E	Timing/Special Directions
aphids (including Russian wheat aphid) brown wheat mite grasshoppers	½ pt/acre	From emergence to flowering, treat when 15-20% of tillers are infested. From flowering to early milk stage, treat when 20% or more of tillers are infested.

wheat midge (orange wheat blossom midge)	½ pt/acre	Treatment is recommended when 75% of the wheat heads have emerged from the boot and when midge adults are found in the crop (1 midge per 4-5 heads). Application timing is critical to ensure good control. If possible, apply in the late afternoon or early evening when temperatures exceed 50°F and wind speed is less than 7 mph.
army cutworms other cutworm species (suppression only)	½ – 1 pt/acre	Control may be reduced under high temperature conditions (greater than 80°F), under dry soil conditions, or if larvae are more than 1/2 inch long. Treat when field counts or crop injury indicates that damaging pest populations are developing or present. A second application of 1 pint/acre may be made for additional control.
cereal leaf beetle	½ – 1 pt/acre	Target application when eggs are near hatching and larvae emerging as monitored by plant inspection.
Restrictions: Do not apply more than 2 applications. Maximum single application rate is 1 pint (0.5 lb ai chlorpyrifos) per acre per season. Do not apply within 14 days of harvest for forage and hay, or within 28 days for grain and straw.		

Mix the required dosage with water and apply in a minimum of 2 gallons per acre finished spray volume. Apply using aerial (fixed wing or helicopter) or power-operated ground spray equipment. For effective coverage of wheat heads using ground application, apply in a minimum of 10 gallons per acre of spray through appropriate nozzles. Higher spray volumes have increased crop protection at the recommended dosage. CPF 4E may also be applied through sprinkler-irrigation systems at recommended broadcast application rates to control listed foliar pests (see directions above).

Restrictions

- Do not make more than 2 applications per crop.
- Do not apply within 28 days of harvest.
- Do not allow livestock to graze or otherwise feed on treated forage within 14 days of application.
- Do not feed straw from treated wheat within 28 days of application.
- Do not apply directly to bodies of water.
- Do not apply product where runoff is likely to occur to aquatic habitats (including lakes, public reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries or other natural waters).
- Do not apply when weather conditions favor drift or runoff from treated areas.

Ground Application

- For ground applications, the distance from treated areas to aquatic habitats (including lakes, public reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries or other natural waters) must be 30 feet or more.
- Do not make ground applications if wind speed is greater than 15 mph.
- Do not apply at spray boom pressures greater than 45 psi.

Aerial Application

- Do not apply by air within 300 feet of aquatic habitats (including lakes, public reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries or other natural waters).
- Do not make aerial applications of CPF 4E when wind speeds exceed 10 mph or when an atmospheric temperature inversion exists.
- Boom length should not exceed 75% of the wing span and release height for aerial applications should be no greater than 10 feet above the crop canopy.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 100°F for extended periods of time. Storage below 20°F may result in formation of crystals. If product crystallizes, store at 50°F to 70°F and agitate to redissolve crystals. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as

indicated below.

PESTICIDE DISPOSAL: Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable containers. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Offer for recycling, if available, or puncture and dispose in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Non-refillable container less than or equal to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Non-refillable container greater than 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over on its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

Spill Cleanup

In Case of Spill: In case of large-scale spillage regarding this product call: CHEMTREC: 1-800-424-9300

Clean-up Procedures: Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing, and wash affected skin areas with soap and water. Keep the spill out of all sewers and open bodies of water,

IMPORTANT INFORMATION - READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of J. Oliver Products, LLC or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of J. Oliver Products, LLC and Seller. To the fullest extent allowed by State law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold J. Oliver Products, LLC and Seller harmless for any claims relating to such factors.

J. Oliver Products, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or J. Oliver Products, LLC, and to the extent consistent with applicable law Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, J. OLIVER PRODUCTS, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, J. Oliver Products, LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF J. OLIVER PRODUCTS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE, AT THE ELECTION OF J. OLIVER PRODUCTS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT, OR COMPENSATION LIMITED TO DAMAGES NOT EXCEEDING THE FAIR MARKET PURCHASE PRICE, AND SHALL NOT INCLUDE INCIDENTAL OR CONSEQUENTIAL DAMAGES.**

J. Oliver Products, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of J. Oliver Products, LLC

Manufactured By:
J. Oliver Products, LLC
3817 Robertson Gin Road
Hernando, MS 38632

EPA Reg. No. 83222-20

EPA Establishment No. 53883-TX-002

44616-MO-001

39578-TX-001

Net Contents: 2.5gallons